RESEARCH

Open Access

Establishment of stable Huh-7 cell lines expressing various hepatitis C virus genotype 3a protein: an in-vitro testing system for novel anti-HCV drugs

Sadia Butt[†], Muhammad Idrees^{*†}, Irshad-ur Rehman[†], Liaqat Ali, Abrar Hussain, Muhammad Ali, Naveed Ahmed, Sana Saleem and Madiha Fayyaz

Abstract

Background: Hepatitis C virus (HCV) infection is the leading cause of chronic hepatitis which progresses to hepatocellular carcinoma (HCC) afflicting > 170 million people worldwide. HCV 3a is the most common genotype (about 70% of all genotypes) circulating in Pakistan. Expression of HCV individual gene of 3a would facilitate therapeutic and vaccines strategies against chronic HCV and liver Cirrhosis. The aim of the present study was the establishment of stable Huh-7 cell lines expressing structural and non structural proteins of HCV Genotype 3a Pakistani isolate obtained from chronic HCV patients.

Methods: Blood samples were obtained from chronic HCV-3a positive patients. HCV individual genes were amplified using PCR with gene specific primers having restriction sites. These gene amplicons were cloned in mammalian expression vector PcDNA3.1+. Huh-7 cell lines were transfected with these constructed plasmids having structural or non-structural HCV genes in confluent cells with lipofectamine. Positive clones were selected with G418 and then confirmed by genome PCR. Subsequently, transcription and expression of the integrated genes were demonstrated by RT-PCR, sequencing and Western blot analysis.

Results: We successfully cloned and express five HCV-3a genes in PcDNA3.1+ mammalian expression vector. Results of western blot and sequencing PCR confirmed the stable expression of these five genes.

Conclusion: The stable cell-lines expressing HCV-3a individual genes would be a useful tool to investigate the role of various HCV proteins on HCV disease outcome and testing of new therapeutic strategies against HCV.

Background

Hepatitis C virus (HCV) is an enveloped plus-strand RNA virus of family *Flaviviridae* [1,2]. HCV is a major leading cause of chronic liver disease [3]. An estimated 170-200 million persons worldwide are infected with HCV [4-6]. Studies on virus replication and pathogenesis having difficulties due to the unavailability of consistent and efficient cell culture systems, even though increasing knowledge of genome structure and individual viral proteins [7]. The HCV genome is approximately 9.6 kb in

length and consists of a single open reading frame (ORF) encoding a polyprotein of about 3,000 amino acids and un-translated regions (UTRs) located at the 5' and 3' terminus of the genome [8,9].

At the 5' end HCV genome there are structural genes; the nucleocapsid region core (C), and the envelope regions (E1 and E2). The 5' UTR and C are conserved regions, while the envelope domain E2/NS1 encloses the hyper variable region [10,11]. After the C gene towards the 3' end, are six non-structural regions (NS2, NS3, NS4A, NS4B, NS5a & NS5B) [7,12]. Viral proteins included in various immunoassays and in the recombinant immuno-blot assay are presented below their corresponding genes [13]. HCV does not integrate into the host genome as it

* Correspondence: idreeskhan@cemb.edu.pk

† Contributed equally

Molecular Virology Laboratory, National Centre of Excellence in Molecular Biology, 87-West Canal Bank Road, Thokar Niaz Baig, Lahore-53700, University of the Punjab, Lahore, Pakistan

does not replicate via a DNA intermediate. Even if the *in-vitro* HCV replication remains a challenge, the chimpanzee is the only important experimental animal model [13]. At the 5'UTR, an internal ribosome entry site (IRES) is located where in a cap-independent manner, viral proteins are expressed. There are 10 viral proteins: core; envelope protein 1 (E1) and E2 are structural proteins that constitute the virion; a small protein that is essential for protein assembly [10,11,14] and six non structural proteins (NS2, NS3, NS4A/B and NS5A/B).

The core protein of HCV forms the nucleocapsid of the virus. It binds with RNA and also interacts with numerous cellular proteins. Various host cell functions such as gene transcription, lipid metabolism, apoptosis and certain signaling pathways are also reported to have interaction with core protein [15]. The associations of core protein with the induction of steatosis and HCC have also been reported [16]. HCV core Ag proved to be useful for performing HCV RNA measurement among dialysis patients in routine laboratories without the need for special equipment or training [17].

E1 protein is associated with the membrane fraction [18]. A direct role for the C-terminal domain in E1 membrane association was identified in the soluble phase by the truncated mutant E1t [19]. The HCV E1 protein having good specificity and could be used in the diagnosis of HCV infection [20] can become useful tools in anti-HCV vaccine research [21]. The NS2 protein is a 23-kDa hydrophobic transmembrane, anchored to the endoplasmic reticulum (ER) [22,23] its function is reliant on the microsomal membranes occurrence, but the function of the NS2 protein in cells is still very poorly understood [24,25]. It has been found that the HCV NS2 protein inhibits cell proliferation and induces cell cycle arrest in the S-phase in mammalian cells through down-regulation of cyclin A expression [24]. Nonstructural protein 4A (NS4A) is a multifunctional protein with 54 amino acid residues. It acts as a cofactor of NS3 serine protease and plays an essential role in the NS4A-dependent cleavage at the NS3-NS4A and NS4B-NS5A junctions [26,27]. Both NS4A and NS4B proteins were previously demonstrated to suppress translation in culture cells [28,29]. HCV NS4B is a highly hydrophobic, localized to the endoplasmic reticulum (ER) and induces a pattern of cytoplasmic foci positive for markers of the ER through four transmembrane segments [30]. NS4B is also a helper factor for the HCV RNA dependent RNA polymerase suggested by the mutagenesis studies of the nucleotide binding motif of NS4B [31]. The involvement of HCV NS4B in IFN- α resistance was also reported by some groups [32,33]. However no such study is available on the construction of these expressions vectors from Pakistan where the rate of HCV is 8-10% in general population and novel and chief drugs are required to treat so huge number of cases.

Therefore, in this study, we have constructed five expression vectors encoding structural (core and envelope1) and nonstructural (NS2, NS4A, NS4B) genes from local HCV isolates and checked their stable expression in Huh-7 cell line. These expression vectors have the potential to be use for testing of new developed drugs in cell culture system.

Methods

Sample collection

Chronic HCV infected with Genotype 3a positive samples were obtained from Division of Molecular Virology and Molecular Diagnostics, National Centre of Excellence in Molecular Biology (CEMB), Lahore, Pakistan. HCV genotyping was carried out on positive HCV PCR samples using type specific HCV genotyping methods as described previously [34,35].

Construction of plasmid (HCV genes in mammalian expression vector PcDNA3.1+)

From the HCV positive serum with 3a genotype, RNA was extracted using Gentra RNA isolation kit (Gentra System Pennsylvania, USA) and individual gene is reverse transcribed using M-MLV (Invitrogen Life technologies, CA). HCV reference sequence of NZL1 # D17763 was used for primer designing on Primer 3 software, restriction sites and kozak sequences were added after restriction analysis on web cutter and neb-cutter primers sequences given in table 1. Each gene is amplified individually and completely. Amplified genes with restriction sites were then cloned in mammalian expression vector PcDNA3.1+ (Invitrogen Life technologies, CA). Each gene constructed plasmid were confirmed through PCR, restriction digestion and sequenced. Individual gene sequence submitted to genbank accession numbers given in table 2.

Cell culture and transfection

Huh-7 cell lines were used maintained in Dulbecco's modified eagle medium (DMEM) supplemented with 100 μ g/ml penicillin; streptomycin and 10% fetal bovine serum (Sigma Aldrich, USA) at 37°C with 5% CO₂. cells were seeded in 24-well (1×10^5 /well) or 6- well (5×10^5 /well) plates and cultured until they became 70-80% confluent. Constructed plasmids about 3-4 μ g of structural (core and E1) and non-structural (NS2, NS4A, NS4B) HCV genes were transfected in confluent cells with lipofectamine (Invitrogen Life technologies, CA) after 6-8 hrs. of transfection media (with lipofectamine and plasmid) was changed.

Isolation of RNA

RNA was isolated using Gentra Kit and reverse transcribed to cDNA with reverse primer and specific genes

Table 1 indicating HCV Gene and polyprotein sequences submitted in Genbank and their Accession Numbers

No.	HCV Gene and polyprotein	Sequences	Accession Numbers
1.	Core	Hepatitis C virus isolate PKIS-1 polyprotein gene, partial cds.	FJ851546.2
		Hepatitis C virus isolate PKIS-2 core polyprotein gene, partial cds.	HQ323687
		Hepatitis C virus isolate PK-1 complete genome.	GU294484.1
2.	Envelope 1	Hepatitis C virus genotype 3a isolate PKIS-2 e1 complete polyprotein gene.	HQ433527
		Hepatitis C virus isolate PK-1 complete genome	GU294484.1
3.	Non-Structural 2	HCV genotype 3a Non-Structural2 NS2 region of Pakistani isolate.	FJ865505
		Hepatitis C virus clone 3a nonstructural protein 2 Pakistani isolate PKIS-2 polyprotein.	HQ822055
		Hepatitis C virus isolate PK-1 complete genome	GU294484.1
4.	Non-structural 4a	Hepatitis C virus isolate PKIS-1 non structural 4a polyprotein gene, partial cds.	HQ822054
		Hepatitis C virus isolate PK-1 complete genome.	GU294484.1
5.	Non-structural 4b	Hepatitis C virus isolate PK1 non-structural protein NS4b gene, partial cds.	GQ325251
		Hepatitis C virus isolate PKIS-2 non-structural protein NS4b gene, partial cds.	HQ323685
		Hepatitis C virus genotype 3a PKIS-3 non-structural protein NS4b.	HQ433528
		Hepatitis C virus genotype 3a isolate PKIS-4 non-structural protein 4b.	HQ616144
		Hepatitis C virus isolate PK-1 complete genome	GU294484.1

were amplified with gene-specific primers for mRNA confirmation.

Proteins extraction and Immuno-blot (Western blotting)

Cells were lysed and protein was extracted after 72 hrs. after transfection and for single stable clones after 3 weeks in PLB (150 mM6/29/2011 NaCl, 1M Tris-Cl pH 7.4, 5 mM EDTA, 1% Triton X-100) proteinase inhibitor and 1 mM PMSE, kept on ice for 15 min. 80-100 µg of total protein were loaded in each well on 10-12.5% SDS-PAGE gels and electrophoretically blotted onto a Hybond-C extra nitrocellulose membrane semi-dry blotting apparatus (Bio-Rad). The membrane was blocked for 1 hour with a 5% milk solution in Phosphate Buffered Saline-0.05% Tween (PBS-T), washed three times with 50 ml of PBS-T. A mixture of primary antibodies for structural genes like core (sc-57800), E1 (sc-65459) and non structural gene like NS4A (sc-52415), NS4B

(sc-65457) was added, each at a concentration of 1:500-1:800 in 5 ml of PBS-T. After incubating at room temperature for 1 hour, the membrane was washed 3 times with PBS-T. A secondary antibody, rabbit anti-mouse IgG, conjugated to alkaline phosphatase (Sigma), was added at a dilution of 1/1000 in PBS-T, incubated at room temperature for one hour. The membrane was washed for three times with PBS-T. Substrate tablet (NBT/BCIP) was dissolved in 1XPBS and blot was incubated for 15-30 min.

Generation of stable cell lines of structural and non structural proteins

After 72 hrs of transfection, cells were given selection with G418 initially with 400 µg/ml for selecting stable clones than after 14 days were given 200 µg/ml. The medium was changed after every 72 hours day. Colonies of G418 resistant cells were selected and grown further

Table 2 List of primers of each individual gene of HCV genotype 3a, Restriction sites worked successfully, Nucleotide position in full length sequence reference sequence of NZL1 was used and number of nucleotides in each amplified region

No.	Genes	Primer seq. 5'-3'	Restriction site	No. of Nucleotides (amplified region)
1.	CORE-IS	ATGAGCACACTTCTAAACCTCA	<i>Hind III</i>	573
2.	CORE-IAS	ACTGGCTGCTGGATGAATTAAGC	<i>EcoRI</i>	
3.	E1-IS	CTAGAGTGGCGGAATACGTCTG	<i>Hind III</i>	
4.	E1-IAS	GGCGACCCCTGAGAACATAACC	<i>EcoRI</i>	576
5.	NS2-IS	CTTTGGTCCCTAGCATTGC	<i>Hind III</i>	
6.	NS2-IAS	CCTCACGGCCTAATCGTGC	<i>EcoRI</i>	642
7.	NS4A-IS	AGCACCTGGGTGTTGCTC	<i>Hind III</i>	
8.	NS4A-IAS	GCACTCCTCCATCTCATCGT	<i>EcoRI</i>	168
9.	NS4B-IS	TCACAAGCTGCCCCATATATCG	<i>Hind III</i>	
10.	NS4B-IAS	GCTACAAGGGCTTGGGTAGTC	<i>XbaI</i>	783

and confirmed with PCR, western blotting and sequencing.

Results

Figure 1 (a & b) showing amplified structural (core and envelope1) and nonstructural (NS2, NS4A, NS4B) genes of the exact sizes. These bands were confirmed by sequencing and only the sequence confirmed genes were further used in next experiment leading to the development of expression vectors. The genes were then cloned in mammalian expression vector pcDNA 3.1+. The successful clones of these genes in PcDNA3.1+ vector were confirmed using restriction digestion analysis. The results of restriction digestions are shown in figure 2. This vector has a CMV promoter which represents an effective mean to transduce eukaryotic cells for transient and stable expression studies. The cloned genes were sequenced in both direction and the consensus sequence was matched to HCV genotype 3a sequence when blast was done with other HCV sequences in GenBank data base.

The expression vector was then linearized and transfected into Huh7 cells by lipofectamine. Twenty-four hours post transfection, selection was applied to the transfected cells by growing them in the presence of 1 mg of G418/ml. About 80% of cells did not develop resistance to the selecting agent, but in the long run it was possible to identify G418-resistant cell clones, which were picked after four weeks of culture and grown as individual cell lines. Once the clones had been isolated and individually grown as cell lines, the concentration of neomycin was decreased to 500 µg/ml. The individual cell lines showed some variability in growth rate.

To check expression of various HCV individual proteins produced from corresponding replicon clones, we performed Western blot analyses with protein extracts

of transfected Huh-7 cells. Figure 3 showing the western blot results of structural and non structural proteins. The Western blot analysis identified specific bands of the expected electrophoretic mobility. B-Actin was used as a positive control. Antibodies of NS2 are not available so it was proceed the same way that was confirmed by sequence analysis and RT-PCR confirmed it. The expression of these individual genes were confirmed by RT PCR and sequencing. All the sequences were submitted to Genbank data base. Table 1 indicating HCV Gene, polyprotein sequences submitted to Genbank data base and their assigned Accession Numbers. Table 2 shows the list of primers of each individual gene of HCV genotype 3a, restriction sites worked successfully, Nucleotide position in full length reference sequence of NZL1 was used and number of nucleotides in each amplified region.

Discussion

Despite vigorous host immune response, 20% of those infected with chronic HCV will eventually lead to HCC [36]. The socio-economic burden of HCV infection globally is striking with an urgent necessity to have better information of viral pathogenesis in order to develop new anti-HCV strategies.

To test novel drugs for its inhibitory action, an efficient culture system is required for the amplification of virus. To date an efficient and reliable culture system is not available to amplify HCV [2] and this limitation prevents the elaboration of reliable infection assays. Several models based on the self-replication of engineered mini-genomes in cell cultures, has been established for HCV replication in other regions of the world [7,37]. The HCV stable cell lines may be very useful in the study of HCV genomic replication in that part of the world

Figure 1 a) Showing the amplified genes of Core (573), Envelope 1(576), Non-structural 2 (NS2, 642 bp) and Non Structural 4a (NS4A, 168 bp), b) lanes 2-5 (left to right) showing the complete amplified region of 783 bp of Non Structural 4b gene. From the HCV positive serum with 3a genotype, RNA was extracted and individual gene was reverse transcribed using M-MLV. HCV reference sequence of NZL1 # D17763 was used for primer designing on Primer 3 software, restriction sites and kozak sequences were added after restriction analysis on web cutter and neb-cutter primers sequences. Each entire gene was amplified individually. Amplified genes with restriction sites were then cloned in mammalian expression vector PcDNA3.1+.

where other HCV genotypes exist. As HCV genotype 3a is the predominant genotype circulating in Pakistan [34,38], therefore, new approaches based on this local HCV genotype 3a are needed on urgent basis to study HCV assembly and infection to design HCV cell entry inhibitors and further to study the humoral immune response against HCV. Therefore, we have developed cell-culture based systems stably expressing two structural and three non-structural HCV individual genes described in the current study.

To the best of our knowledge no cell culture based system has yet been developed to propagate the replication expression of HCV 3a genes of Pakistani chronic isolates in cultured mammalian cells. Because the existing replicon was generated using genotype 1b HCV RNA, the present replicon system may not be used to detect responses that are genotype and subtype-dependent. Therefore this study was initiated to establish stable cell lines expressing proteins of Pakistani HCV genotype 3a isolates. The establishment of HCV genotype 3a cell lines stably expressing structural and non structural proteins is an instrumental in the further study of HCV replication and viral-host interaction of genotype 3a. Viral and cellular factors required for HCV replication will be defined by cutting edge gene and micro-array, proteomics, protein-protein interactions methodologies. Further investigation on these stable cell lines must have direct impact on HCV disease outcome and new therapeutic strategies will be designed.

Conclusion

In summary, we were able to develop vectors stably expressing HCV individual proteins such as core, envelope1 (Structural), NS2, NS4A and NS4B (Non-structural). The stable cell line expressing individual HCV gene would be useful in identifying the role of most important genes in HCC and fibrosis development and studying the mechanisms of each gene in HCV replication. Obviously, novel therapeutic strategies are required on urgent basis as the health costs for HCV-infected people are predicted to spiral dramatically in the next decade. Further investigation on these stable cell lines must have direct impact on HCV disease outcome and new therapeutic strategies will be designed. This system with genes from HCV-3a strain can be used for comparison studies with other strain-derived systems developed in other areas for the analysis of the effects of anti-HCV drugs.

Acknowledgements

We thank all the clinicians and patients for their cooperation in the study.

Authors' contributions

SB and IR reviewed the literature, conducted all the experiments and wrote the manuscript. MI guided conducting the whole experiment and edited the manuscript. LA, MA, AH, BR, SS, NA, helped SB & IR in literature review. All the authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 5 April 2011 Accepted: 28 June 2011 Published: 28 June 2011

References

- Ogata N, Alter HJ, Miller RH, Purcell RH: **Nucleotide sequence and mutation rate of the H strain of hepatitis C virus.** *Proc Natl Acad Sci* 1991, **88**:3392-3396.
- Lindenbach BD, Thiel HJ, Rice CM: **Flaviviridae: the viruses and their replication.** In *Fields virology. Volume 5.* 5 edition. Edited by: Fields BN,

- Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE. Lippincott, Williams and Wilkins, Philadelphia, PA; 2007:1101-1152.
3. Poynard T, Ratziu V, Benhamou Y, Opolon P, Cacoub P, Bedossa P: **Natural history of HCV infection.** *Best Pract Res Clin Gastroenterol* 2000, **14**:211-228.
4. Kato T, Furusaka A, Miyamoto M, Date T, Yasui K, Hiramoto J, Nagayama K, Tanaka T, Wakita T: **Sequence analysis of hepatitis C virus isolated from a fulminant hepatitis patient.** *J Med Virol* 2001, **64**(3):334-9.
5. Shepard CW, Finelli L, Alter MJ: **Global epidemiology of hepatitis C virus infection.** *Lancet Infect. Dis* 2005, **5**:558-567.
6. Chen SL, Morgan TR: **The natural history of hepatitis C virus (HCV) infection.** *Int J Med Sci* 2006, **3**:47-52.
7. Lohmann V, Körner F, Koch JO, Herian U, Theilmann L, Bartenschlager R: **Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line.** *Science* 1999, **285**:110-113.
8. Lindenbach BD, Rice CM: **Flaviviridae: the viruses and their replication.** *Fields Virology* Philadelphia: Lippincott Williams & Wilkins; 2001.
9. Lemon SM, Walker CM, Alter MJ, Yi M: **Fields Virology.** In *Hepatitis C virus*. Edited by: Knipe DM, Howley PM. Lippincott Williams and Wilkins, Philadelphia; 2007:1253-1304.
10. Jones CT, Murray CL, Eastman DK, Tassello J, Rice CM: **Hepatitis C virus p7 and NS2 proteins are essential for production of infectious virus.** *J Virol* 2007, **81**:8374-8383.
11. Steinmann E, Penin F, Kallis S, Patel AH, Bartenschlager R, Pietschmanns T: **Hepatitis C virus p7 protein is necessary for assembly and release of infectious virions.** *PLoS Pathog* 2007, **3**:e103.
12. Gosert R, Egger D, Lohmann V, Bartenschlager R, Blum HE, Bienz K, Moradpour D: **Identification of the hepatitis C virus RNA replication complex in Huh-7 cells harboring subgenomic replicons.** *J Virol* 2003, **77**:5487-5492.
13. Dienstag Dienstag JL, Isselbacher KJ: **Acute viral hepatitis.** In *Iskasper DL Harrison's Principles of Internal Medicine. Volume II.* 16 edition. New York: McGraw-Hill; 2005:1829.
14. Griffin SD, Beales LP, Clarke DS, Worsfold O, Evans SD, Jaeger J, Harris MP, Rowlands DJ: **The p7 protein of hepatitis C virus forms an ion channel that is blocked by the antiviral drug, amantadine.** *FEBS Lett* 2003, **535**:34-38.
15. Tellinghuisen TL, Rice CM: **Interaction between hepatitis C virus proteins and host cell factors.** *Curr Opin Microbiol* 2002, **5**:419-27.
16. Lerat H, Honda M, Beard MR, Loesch K, Sun J, et al: **Steatosis and liver cancer in transgenic mice expressing the structural and nonstructural proteins of hepatitis C virus.** *Gastroenterol* 2002, **122**:352-65.
17. Fabrizi F, Lunghi G, Aucella F, Mangano S, Barbisoni F, Bisegna S, Vigilante D, Limido A, Martin P: **Novel assay using total hepatitis C virus (HCV) core antigen quantification for diagnosis of HCV infection in dialysis patients.** *J Clin Microbiol* 2005, **43**(1):414-20.
18. Susanne N, Karin S, Christoph K: **Expression of Semliki Forest Virus E1 Protein in Escherichia coli.** *J Bio Chem* 2001, **276**(18):15453-15457.
19. Ciccaglione AR, Marcantonio C, Costantino A, Equestre M, Geraci A, Rapicetta M: **Expression and membrane association of hepatitis C virus envelope 1 protein.** *Virus Genes* 2000, **21**(3):223-6.
20. Gao J, Tao Q, Ma D: **Prokaryotic expression of hepatitis C virus envelope 1 gene and application of the expressed product.** *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 2001, **15**(1):20-3.
21. Liu J, Zhu LX, Kong YY, Li GD, Wang Y: **Purification and application of C-terminally truncated hepatitis C virus E1 proteins expressed in Escherichia coli.** *World J Gastroenterol* 2005, **11**(4):503-507.
22. Santolini E, Pacini L, Fipaldini C, Migliaccio G, Monica N: **The NS2 protein of hepatitis C virus is a transmembrane polypeptide.** *J Virol* 1995, **69**:7461-7471.
23. Yamaga AK, Ou JH: **Membrane topology of the hepatitis C virus NS2 protein.** *J Biol Chem* 2002, **277**:33228-33234.
24. Yang XJ, Liu J, Ye L, Liao QJ, Wu JG, Gao JR, She YL, Wu ZH, Ye LB: **HCV NS2 protein inhibits cell proliferation and induces cell cycle arrest in the S-phase in mammalian cells through down-regulation of cyclin A expression.** *Vir Res* 2006, **121**(2):134-143.
25. Bussche A, Machida R, Li K, Loevinsohn G, Khander A, Wang J, Wakita T, Wands JR, Li J: **Hepatitis C virus NS2 protein triggers endoplasmic reticulum stress and suppresses its own viral replication.** *J Hepatol* 2010, **53**(5):797-804.
26. Failla C, Tomei L, De Francesco R: **Both NS3 and NS4A are required for proteolytic processing of hepatitis C virus nonstructural proteins.** *J Virol* 1994, **68**:3753-3760.
27. Lin C, Pragai BM, Grakoui A, Xu J, Rice CM: **Hepatitis C virus NS3 serine proteinase: trans-cleavage requirements and processing kinetics.** *J Virol* 1994, **68**:8147-8157.
28. Florese RH, Nagano-Fujii M, Iwanaga Y, Hidajat R, Hotta H: **Inhibition of protein synthesis by the nonstructural proteins NS4A and NS4B of hepatitis C virus.** *Virus Res* 2002, **90**:119-131.
29. Kato J, Kato NY, Yoshida H, Nita SKO, Shiratori Y, Omata M: **Hepatitis C virus NS4A and NS4B proteins suppress translation in vivo.** *J Med Virol* 2002, **66**:187-199.
30. Lundin M, Monne M, Widell A, VonHeijne G, Persson MA: **Topology of the membrane-associated hepatitis C virus protein NS4B.** *J Virol* 2003, **77**(9):5428-5438.
31. Einav S, Elazar M, Danieli T, Glenn JS: **A nucleotide binding motif in hepatitis C virus (HCV) NS4B mediates HCV RNA replication.** *J Virol* 2004, **78**:11288-11295.
32. Munoz-Jordan JL, Laurent-Rolle M, Ashour J, Martinez-Sobrido L, Ashok M, Lipkin WJ, Garcia-Sastre A: **Inhibition of alpha/beta interferon signaling by the NS4B protein of flaviviruses.** *J Virol* 2005, **79**(13):8004-8013.
33. Jing Xu, Shufeng Liu, Yihui Xu, Po Tien, Guangxia Gao: **Identification of the nonstructural protein 4B of hepatitis C virus as a factor that inhibits the antiviral activity of interferon-alpha.** *Elsevier Vir Res* 2009, **141**:55-62.
34. Idrees M, Riazuddin S: **Frequency distribution of hepatitis C virus genotypes in different geographical regions of Pakistan and their possible routes of transmission.** *BMC Infect Dis* 2008, **8**:69.
35. Ohno T, Mizokami M, Wu RR, Saleh GM, Ohba KI, Orito E, Mukaide M, Williams R, Lau JYN: **New hepatitis C virus (HCV) genotyping system that allows for identification of HCV genotypes 1a, 1b, 2a, 2b, 3a, 3b, 4, 5a, and 6a.** *J Clin Microbiol* 1997, **35**:201-207.
36. Sheehy P, Scallan M, Kenny-Walsh S, Shanahan F, Fanning LJ: **A strategy for obtaining near full-length HCV cDNA clones (assemblons) by assembly PCR.** *J Virol* 2005, **123**(2):115-124.
37. Blight KJ, Grakoui A, Hanson HL, Rice CM, Ou J-HJ: **The molecular biology of hepatitis C virus.** *Hepatitis viruses* Boston: Kluwer Academic Publishers; 2002, 81-108.
38. Butt S, Idrees M, Akbar H, Rehman I, Awan Z, Afzal S, Hssain A, Shahid M, Manzoor S, Rafique S: **The changing epidemiology pattern and frequency distribution of hepatitis C virus in Pakistan.** *Infect Genet Evol* 2010, **10**(5):595-600.

doi:10.1186/1479-0556-9-12

Cite this article as: Butt et al.: Establishment of stable Huh-7 cell lines expressing various hepatitis C virus genotype 3a protein: an in-vitro testing system for novel anti-HCV drugs. *Genetic Vaccines and Therapy* 2011 9:12.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit